




Λ_c baryon production in Au+Au collisions at

$$\sqrt{s_{NN}} = 200 \text{ GeV}$$

Miroslav Simko, for the STAR Collaboration

Nuclear Physics Institute of the Czech Academy of Sciences,

Faculty of Nuclear Sciences and Physical Engineering of Czech Technical University in Prague



2015 KOBE JAPAN

Baryon-to-meson yield ratios (p/π , Λ/K^0) are observed to be significantly enhanced in central heavy-ion collisions than those in peripheral heavy-ion collisions and p+p collisions at RHIC and LHC. This effect can be explained by taking into account coalescence hadronization that recombine thermalized light and strange quarks in heavy-ion collisions as opposed to fragmentation hadronization in elementary nucleon-nucleon collisions. Several model calculations suggest that coalescence hadronization between charm quarks and light quarks will lead to an enhancement in the Λ_c/D^0 ratio. The possible Λ_c/D^0 enhancement in heavy-ion collisions will introduce additional suppression for charm decay electrons due to smaller semi-leptonic decay branching ratios of Λ_c , which could lead to a different interpretation of the heavy flavor decay electron results.

Λ_c baryons have an extremely small lifetime ($c\tau \sim 60 \mu\text{m}$) and have not been measured in heavy-ion collisions yet. The newly installed STAR Heavy Flavor Tracker (HFT) has shown high efficiency and excellent pointing resolution that can facilitate Λ_c reconstruction in heavy-ion collisions. We report studies of reconstruction of Λ_c baryons via three body hadronic decays into p K and π , using year 2014 Au+Au data at $\sqrt{s_{NN}} = 200 \text{ GeV}$.

Λ_c baryon

- Ratios of baryons/mesons (such as p/π , Λ/K^0) in heavy-ion collisions significantly enhanced compared to p+p collisions – due to quark coalescence.
- Similar enhancement expected in the ratio of Λ_c/D^0 .
- This would bring insight into coalescence hadronization of charm quarks.
- The enhancement of Λ_c/D^0 is one of the signatures of the formation of the strongly coupled Quark-Gluon Plasma.
- Challenging to measure, $c\tau \sim 60 \mu\text{m}$.
- Λ_c have never been observed in high energy heavy-ion collisions.

Λ_c branching ratios for three body decay:

- $\Lambda_c^+ \rightarrow p K^- \pi^+$ ($5.0 \pm 1.3\%$)
- $\Lambda_c^+ \rightarrow p K^*$ ($1.6 \pm 0.5\%$)
- $\Lambda_c^+ \rightarrow \Lambda(1520) \pi^+$ ($1.8 \pm 0.6\%$)
- $\Lambda_c^+ \rightarrow K^- \Delta^{++}$ ($0.86 \pm 0.3\%$)
- Nonresonant ($2.6 \pm 0.8\%$)

[Particle Data Group, Chin. Phys. C38 (2014) 090001]




Figure 1: Measurement of Λ_c/K_S^0 in heavy-ion and p+p collisions. Strong enhancement is observed in intermediate p_T [J. Phys. Conf. Ser. 50 (2006) 192].

Comparison: Simulation and Background from data

- p K π channel – triplets of Λ_c candidates.
- Simulated Λ_c compared to background from 200 GeV Au+Au data.
- Background was selected from p K π triplets with wrong sign combinations of the daughters.
- The same cuts were applied on background and Λ_c from simulation.




Figure 5: Distribution of variables: Decay length, daughters p_T , $\cos(\theta)$, distance of pairs vertices, and daughters DCA for signal simulation and background from data. The background is scaled to show the difference in shape.

Detector layout




Figure 2: The STAR experiment and subdetectors used in the Λ_c analysis.

Kinematic and topological variables for Λ_c analysis

- Decay length:** distance of secondary vertex to the primary vertex
- θ :** angle between Λ_c momentum and position relative to primary vertex
- Daughter pair DCA:** distance of closest approach (DCA) between the daughter pairs
- Distance between vertices of daughter pairs**
- Daughter p_T**




Figure 3: Variables used in topological cuts

Λ_c signal simulation

- Data driven Monte Carlo simulation was used:
- Λ_c decayed in Pythia.
- p_T and positions of the daughter particles were smeared, according to resolution observed in data.
- Λ_c were generated with flat rapidity distribution within $|y| < 1$.
- The p_T distribution was obtained from the D^0 minimum bias spectrum from 0–80% centrality Au+Au collisions [Phys. Rev. Lett. 113 (2014) 142301].
- Fragmentation ratio 0.275 between Λ_c and D^0 was taken from ZEUS data [Eur. Phys. J. C44 (2005) 351].
- Scaling of Λ_c production with the number of binary collisions was used.




Figure 4: D^0 spectrum fitted by Levy function (used as p_T distribution for Λ_c)

Run 2016 and future upgrades

- STAR has recorded 1.2 B minimum bias events in Au+Au collisions at $\sqrt{s_{NN}} = 200 \text{ GeV}$ in the year 2014 and plans to take 2 B more in 2016.
- The inner HFT pixel layer will have reduced material budget thanks to changed cable material of the sensors from copper to aluminum.
- The HFT will have higher efficiency because of replacing non-working sensors.
- This increases the pointing and p_T resolution of STAR and increases the efficiency of tracking with HFT.




Figure 6: Comparison of the DCA resolution in XY plane with aluminum (left) and copper cables (right).

Conclusion and outlook

- Λ_c measurements are ongoing with the new Heavy Flavor Tracker at STAR.
- Λ_c reconstruction efficiency will increase in run 2016 thanks to reduced mass of the HFT infrastructure and reduced number of dead sensors.
- With the statistics of the future runs, STAR could measure Λ_c baryons for the first time thanks to excellent pointing resolution of the HFT.

Acknowledgement: This work was supported by Grant Agency of the Czech Technical University in Prague, grant No. SGS13/215/OHK4/35/14 and by Grant Agency of the Czech Republic, grant No. 13-20841S.

